

STERIMAX INC.

Suite 1, 2735 Matheson Blvd. E., Mississauga, ON, Canada L4W 4M8 • www.sterimaxinc.com
Tel.: 905-890-0661 • 800-881-3550 • Fax: 905-890-0508 • 877-546-7667

New Packaging for Apresoline 20 mg/amp (DIN 00723754)

In our continued efforts to improve our products, we are pleased to announce updated packaging for **Apresoline 20 mg/amp (DIN 00723754)**.

This exciting new packaging better conforms to the **CSHP** and **CSA** “Guidelines for Drug Packaging and Labeling for Manufacturers”.

To help you better understand the changes made, we have outlined the critical changes on the following pages. A number of other smaller ‘cosmetic’ changes were made as well, but not detailed below.

Please note that the product remains the same; only the printed packaging has changed. A sample of the new packaging can be provided to you upon request.

We anticipate the product to be released July 2010; however this is an estimation based on when the inventory level of product available in the current packaging is expected to be depleted. The date may change based on inventory levels.

If you have any questions regarding this update, please feel free to contact SteriMax customer service at 905-890-0661.

STERIMAX INC.

Suite 1, 2735 Matheson Blvd. E., Mississauga, ON, Canada L4W 4M8 • www.sterimaxinc.com
Tel.: 905-890-0661 • 800-881-3550 • Fax: 905-890-0508 • 877-546-7667

Current Carton:

Apresoline®
hydrALAZINE hydrochloride
for injection

20 mg / ampoule

Sterile • For intravenous use.
Store powder below 30°C,
protect from heat and light.

Antihypertensive


10 ampoules

STERIMAX INC.

DIN 00723754

Reconstitution: I.V.: Add 1 mL of Sterile Water for Injection, gives approximately 19.7 mg/mL. Use immediately. I.V. infusion: Reconstitute as above, further dilute with physiological saline (or other recommended I.V. solution). Use within 24 hours.

Adult Dose: Patient should be hospitalized. Use under strict medical supervision. Initially 5 to 10 mg by slow I.V., monitor blood pressure and heart rate every 5 minutes. Dose may be repeated at 20 to 30 minute intervals. May also be given by I.V. infusion, initially at rate of 200 to 300 µg/minute. Maintenance: 50 to 150 µg/minute. See package insert for detailed directions for use. Product monograph supplied on request.



Carton dimension changed from 171 x 78 x 18 mm to 96.84 x 63.5 x 46.04 mm.

Changed inner plastic tray to cardboard tray.

New Carton:

Apresoline®
hydrALAZINE hydrochloride
for injection

20 mg / ampoule

Sterile • For intravenous use.
Store powder below 30°C,
protect from heat and light.

Antihypertensive

10 ampoules

STERIMAX INC.

DIN 00723754

Reconstitution: I.V.: Add 1 mL of Sterile Water for Injection, gives approximately 19.7 mg/mL. Use immediately. I.V. infusion: Reconstitute as above, further dilute with physiological saline (or other recommended I.V. solution). Use within 24 hours.

Adult Dose: Patient should be hospitalized. Use under strict medical supervision. Initially 5 to 10 mg by slow I.V., monitor blood pressure and heart rate every 5 minutes. Dose may be repeated at 20 to 30 minute intervals. May also be given by I.V. infusion, initially at rate of 200 to 300 µg/minute. Maintenance: 50 to 150 µg/minute. See package insert for detailed directions for use. Product monograph supplied on request.




© Reg. TM Novartis
L.E. User, SteriMax Inc.
Mississauga, Ontario
L4W 4M8

ST-CC00500.01

STERIMAX INC.

Suite 1, 2735 Matheson Blvd. E., Mississauga, ON, Canada L4W 4M8 • www.sterimaxinc.com
Tel.: 905-890-0661 • 800-881-3550 • Fax: 905-890-0508 • 877-546-7667

Current Label:



Added GS1 Databar code.

New Label:

